



Trigonometric Functions 6B


1 a




b



c




2 a



b 2 solutions

3 a



b You can see that the graphs of  $y = \sec \theta$  and  $y = -\cos \theta$  do not meet, so  $\sec \theta = -\cos \theta$  has no solutions.

The same result can be found algebraically


$$\sec \theta = -\cos \theta$$

$$\Rightarrow \frac{1}{\cos \theta} = -\cos \theta$$

$$\Rightarrow \cos^2 \theta = -1$$

There are no solutions of this equation for real  $\theta$ .



4 a



b The curves meet at the maxima and minima of  $y = \sin 2\theta$ , and on the  $\theta$ -axis at odd integer multiples of  $90^\circ$ .

In the interval  $0 \leq \theta \leq 360^\circ$  there are 6 intersections. So there are 6 solutions of  $\cot \theta = \sin 2\theta$  in the interval  $0 \leq \theta \leq 360^\circ$

5 a



b  $y = \cot(\theta + 90^\circ)$  is a reflection in the  $\theta$ -axis of  $y = \tan \theta$ , so  
 $\cot(\theta + 90^\circ) = -\tan \theta$


6 a i The graph of  $y = \tan\left(\theta + \frac{\pi}{2}\right)$  is the same as that of  $y = \tan \theta$  translated by the vector  $\begin{pmatrix} -\frac{\pi}{2} \\ 0 \end{pmatrix}$ , i.e. by  $\frac{\pi}{2}$  to the left.

ii The graph of  $y = \cot(-\theta)$  is the same as that of  $y = \cot \theta$  reflected in the  $y$ -axis.

iii The graph of  $y = \operatorname{cosec}\left(\theta + \frac{\pi}{4}\right)$  is the same as that of  $y = \operatorname{cosec} \theta$  translated by the vector  $\begin{pmatrix} -\frac{\pi}{4} \\ 0 \end{pmatrix}$

iv The graph of  $\sec\left(\theta - \frac{\pi}{4}\right)$  is the same as that of  $y = \sec \theta$  translated by the vector  $\begin{pmatrix} \frac{\pi}{4} \\ 0 \end{pmatrix}$


b



(reflection of  $y = \cot \theta$  in the  $y$ -axis)


$$\tan\left(\theta + \frac{\pi}{2}\right) = \cot(-\theta)$$

6 b



$$\operatorname{cosec}\left(\theta + \frac{\pi}{4}\right) = \sec\left(\theta - \frac{\pi}{4}\right)$$


- 7 a A stretch of  $y = \sec \theta$  in the  $\theta$  direction with scale factor  $\frac{1}{2}$   
 Minimum at  $(180^\circ, 1)$   
 Maxima at  $(90^\circ, -1)$  and  $(270^\circ, -1)$   
 It meets the  $y$ -axis at  $(0, 1)$




- b Reflection in  $\theta$ -axis of  $y = \operatorname{cosec} \theta$

Minimum at  $(270^\circ, 1)$

Maximum at  $(90^\circ, -1)$



- c Translation of  $y = \sec \theta$  by the vector  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ , i.e. +1 in the  $y$  direction.  
 It meets  $x$ -axis at  $(180^\circ, 0)$   
 There is a maximum at  $(180^\circ, 0)$   
 It meets the  $y$ -axis at  $(0, 2)$




- d** Translation of  $y = \operatorname{cosec} \theta$  by the vector  $\begin{pmatrix} 30 \\ 0 \end{pmatrix}$

Minimum at  $(120^\circ, 1)$

Maximum at  $(300^\circ, -1)$

It meets the  $y$ -axis at  $(0, -2)$




- 7 e**  $y = 2 \sec(\theta - 60^\circ)$  is  $y = \sec \theta$  translated by the vector  $\begin{pmatrix} 60 \\ 0 \end{pmatrix}$  and then stretched by a scale factor 2 in the  $y$  direction.

Minimum at  $(60^\circ, 2)$

Maximum at  $(240^\circ, -2)$

It meets the  $y$ -axis at  $(0, 4)$




- f**  $y = \operatorname{cosec}(2\theta + 60^\circ)$  is  $y = \operatorname{cosec} q$

translated by the vector  $\begin{pmatrix} -60 \\ 0 \end{pmatrix}$  and then stretched by a scale factor  $\frac{1}{2}$  in the  $\theta$  direction.

Minima at  $(15^\circ, 1), (195^\circ, 1)$


Maxima at  $(105^\circ, -1), (285^\circ, -1)$

It meets the  $y$ -axis at  $(0, 1.155)$



- g**  $y = -\cot 2\theta$  is  $y = \cot \theta$  stretched by a scale factor  $\frac{1}{2}$  in the  $\theta$  direction and then reflected in the  $x$ -axis.

It meets the  $\theta$ -axis at  $(45^\circ, 0)$ ,  $(135^\circ, 0)$ ,  $(225^\circ, 0)$  and  $(315^\circ, 0)$




- h**  $y = 1 - 2\sec \theta = -2\sec \theta + 1$  is  $y = \sec \theta$  stretched by a scale factor 2 in the  $y$  direction, reflected in the  $x$ -axis and then translated by the vector  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$

Minima at  $(180^\circ, 3)$

Maxima at  $(0^\circ, -1)$ ,  $(360^\circ, -1)$

It meets the  $y$ -axis at  $(0, -1)$



- 8 a** The period of  $\sec \theta$  is  $2\pi$  radians  
 $y = \sec 3\theta$  is a stretch of  $y = \sec \theta$  with scale factor  $\frac{1}{3}$  in the  $\theta$  direction.

So the period of  $\sec 3\theta$  is  $\frac{2\pi}{3}$


- b**  $\operatorname{cosec} \theta$  has a period of  $2\pi$   
 $\operatorname{cosec} \frac{1}{2}\theta$  is a stretch of  $\operatorname{cosec} \theta$  in the  $\theta$  direction with scale factor 2.  
So the period of  $\operatorname{cosec} \frac{1}{2}\theta$  is  $4\pi$

- c**  $\cot \theta$  has a period of  $\pi$   
 $2\cot \theta$  is a stretch in the  $y$  direction by scale factor 2. So the periodicity is not affected.

The period of  $2\cot \theta$  is  $\pi$


- d**  $\sec \theta$  has a period of  $2\pi$   
 $\sec(-\theta)$  is a reflection in the  $y$ -axis.  
So the periodicity is not affected.  
The period of  $\sec(-\theta)$  is  $2\pi$

- 9 a**  $y = 3 + 5\operatorname{cosec} \theta$  is  $y = \operatorname{cosec} \theta$  stretched by a scale factor 5 in the  $y$  direction and then translated by the vector  $\begin{pmatrix} 0 \\ 3 \end{pmatrix}$



- b**  $-2 < k < 8$

10 a



b The  $\theta$  coordinates at points at which the gradient is zero are at the maxima and minima. These are  $\theta = -\pi, 0, \pi, 2\pi$

c Minimum value of  $\frac{1}{1+2\sec\theta}$

is where  $1+2\sec\theta$  is a maximum.

So minimum value of  $\frac{1}{1+2\sec\theta}$

is  $\frac{1}{-1} = -1$

The first positive value of  $\theta$  where this occurs is when  $\theta = \pi$

(see diagram)

Maximum value of  $\frac{1}{1+2\sec\theta}$

is where  $1+2\sec\theta$  is a minimum.

So maximum value of  $\frac{1}{1+2\sec\theta}$  is  $\frac{1}{3}$

The first positive value of  $\theta$  where this occurs is when  $\theta = 2\pi$

(see diagram)