

## Trigonometry and modelling 7A

$$\begin{aligned} \mathbf{1\ a\ i}\quad \angle FAB &= \angle CAF + \angle BAC \\ &= (\alpha - \beta) + \beta = \alpha \\ \text{So } \angle FAB &= \alpha \end{aligned}$$

$$\begin{aligned} \mathbf{ii}\quad \angle FAB \text{ and } \angle ABD &\text{ are alternate angles} \\ \text{so } \angle FAB &= \angle ABD \\ \text{so } \angle ABD &= \alpha \\ \angle CBE = 90 - \alpha, \text{ so } \angle ECB &= 90 - (90 - \alpha) = \alpha \end{aligned}$$

$$\begin{aligned} \mathbf{iii}\quad \cos \beta &= \frac{AB}{1} \\ \text{So } AB &= \cos \beta \end{aligned}$$

$$\begin{aligned} \mathbf{iv}\quad \sin \beta &= \frac{BC}{1} \\ \text{So } BC &= \sin \beta \end{aligned}$$

$$\begin{aligned} \mathbf{b\ i}\quad \angle ABD = \alpha, \text{ so } \sin \alpha &= \frac{AD}{AB} \\ \text{As } AB = \cos \beta, \text{ this gives } \sin \alpha &= \frac{AD}{\cos \beta} \\ \text{So } AD &= \sin \alpha \cos \beta \end{aligned}$$

$$\begin{aligned} \mathbf{ii}\quad \cos \alpha &= \frac{BD}{AB} = \frac{BD}{\cos \beta} \\ \text{So } BD &= \cos \alpha \cos \beta \end{aligned}$$

$$\begin{aligned} \mathbf{c\ i}\quad \angle ECB = \alpha, \text{ so } \cos \alpha &= \frac{CE}{BC} \\ \text{As } BC = \sin \beta, \text{ this gives } \cos \alpha &= \frac{CE}{\sin \beta} \\ \text{So } CE &= \cos \alpha \sin \beta \end{aligned}$$


$$\begin{aligned} \mathbf{ii}\quad \sin \alpha &= \frac{BE}{BC} = \frac{BE}{\sin \beta} \\ \text{So } BE &= \sin \alpha \sin \beta \end{aligned}$$

$$\begin{aligned} \mathbf{d\ i}\quad \sin(\alpha - \beta) &= \frac{FC}{1} \\ \text{So } FC &= \sin(\alpha - \beta) \end{aligned}$$

1 d ii  $\cos(\alpha - \beta) = \frac{FA}{1}$

So  $FA = \cos(\alpha - \beta)$

e i The completed diagram should look like this:



$FC + CE = AD$ , so  $FC = AD - CE$   
 $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$

ii  $AF = DB + BE$   
 $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$

2  $\tan(A - B) = \frac{\sin(A - B)}{\cos(A - B)}$   
 $= \frac{\sin A \cos B - \cos A \sin B}{\cos A \cos B + \sin A \sin B}$

Divide the numerator and denominator by  $\cos A \cos B$

$$\tan(A - B) = \frac{\frac{\sin A \cos B}{\cos A \cos B} - \frac{\cos A \sin B}{\cos A \cos B}}{\frac{\cos A \cos B}{\cos A \cos B} + \frac{\sin A \sin B}{\cos A \cos B}} = \frac{\frac{\sin A}{\cos A} - \frac{\sin B}{\cos B}}{1 + \frac{\sin A \sin B}{\cos A \cos B}}$$

$$= \frac{\tan A - \tan B}{1 + \tan A \tan B} \text{ as required}$$

3  $\sin(A + B) = \sin A \cos B + \cos A \sin B$   
 $\sin(P + (-Q)) = \sin P \cos(-Q) + \cos P \sin(-Q)$   
 As  $\cos(-P) = \cos P$  and  $\sin(-P) = -\sin P$ , this gives  
 $\sin(P - Q) \equiv \sin P \cos Q - \cos P \sin Q$

4 Example:  $A = 60^\circ$ ,  $B = 30^\circ$

$$\sin A = \frac{\sqrt{3}}{2}, \sin B = \frac{1}{2}$$

$$\sin(A+B) = 1; \sin A + \sin B = \frac{\sqrt{3}}{2} + \frac{1}{2} \neq 1$$

This proves  $\sin(A+B) = \sin A + \sin B$  is not true for all values.

There will be many values of  $A$  and  $B$  for which the statement is true, e.g.  $A = -30^\circ$  and  $B = +30^\circ$ , and this is the danger of trying to prove a statement by taking particular examples. To prove a statement requires a sound argument; to disprove it only requires one counterexample.

5  $\cos(A-B) \stackrel{?}{=} \cos A \cos B + \sin A \sin B$

Set  $A = \theta$ ,  $B = \theta$

$$\Rightarrow \cos(\theta - \theta) \equiv \cos \theta \cos \theta + \sin \theta \sin \theta$$

$$\Rightarrow \cos 0 \equiv \cos^2 \theta + \sin^2 \theta$$

So  $\cos^2 \theta + \sin^2 \theta \equiv 1$  (since  $\cos 0 = 1$ )

6 a  $\sin(A-B) \stackrel{?}{=} \sin A \cos B - \cos A \sin B$

Set  $A = \frac{\pi}{2}$ ,  $B = \theta$

$$\Rightarrow \sin\left(\frac{\pi}{2} - \theta\right) \equiv \sin \frac{\pi}{2} \cos \theta - \cos \frac{\pi}{2} \sin \theta$$

$$\Rightarrow \sin\left(\frac{\pi}{2} - \theta\right) \equiv \cos \theta$$

since  $\sin \frac{\pi}{2} = 1$ ,  $\cos \frac{\pi}{2} = 0$

b  $\cos(A-B) \stackrel{?}{=} \cos A \cos B + \sin A \sin B$

Set  $A = \frac{\pi}{2}$ ,  $B = \theta$

$$\Rightarrow \cos\left(\frac{\pi}{2} - \theta\right) \equiv \cos \frac{\pi}{2} \cos \theta + \sin \frac{\pi}{2} \sin \theta$$

$$\Rightarrow \cos\left(\frac{\pi}{2} - \theta\right) \equiv \sin \theta$$

since  $\cos \frac{\pi}{2} = 0$ ,  $\sin \frac{\pi}{2} = 1$

7  $\sin\left(x + \frac{\pi}{6}\right) = \sin x \cos \frac{\pi}{6} + \cos x \sin \frac{\pi}{6}$

$$= \frac{\sqrt{3}}{2} \sin x + \frac{1}{2} \cos x$$

$$\begin{aligned}
 8 \quad \cos\left(x + \frac{\pi}{3}\right) &= \cos x \cos \frac{\pi}{3} - \sin x \sin \frac{\pi}{3} \\
 &= \frac{1}{2} \cos x - \frac{\sqrt{3}}{2} \sin x
 \end{aligned}$$

**9 a** Using  $\sin(A+B) \equiv \sin A \cos B + \cos A \sin B$  gives  
 $\sin 15^\circ \cos 20^\circ + \cos 15^\circ \sin 20^\circ \equiv \sin(15^\circ + 20^\circ) \equiv \sin 35^\circ$

**b** Using  $\sin(A-B) \equiv \sin A \cos B - \cos A \sin B$  gives  
 $\sin 58^\circ \cos 23^\circ - \cos 58^\circ \sin 23^\circ \equiv \sin(58^\circ - 23^\circ) \equiv \sin 35^\circ$

**c** Using  $\cos(A+B) \equiv \cos A \cos B - \sin A \sin B$  gives  
 $\cos 130^\circ \cos 80^\circ - \sin 130^\circ \sin 80^\circ \equiv \cos(130^\circ + 80^\circ) \equiv \cos 210^\circ$

**d** Using  $\tan(A-B) \equiv \frac{\tan A - \tan B}{1 + \tan A \tan B}$  gives  
 $\frac{\tan 76^\circ - \tan 45^\circ}{1 + \tan 76^\circ \tan 45^\circ} \equiv \tan(76^\circ - 45^\circ) \equiv \tan 31^\circ$

**e** Using  $\cos(A-B) \equiv \cos A \cos B + \sin A \sin B$  gives  
 $\cos 2\theta \cos \theta + \sin 2\theta \sin \theta \equiv \cos(2\theta - \theta) \equiv \cos \theta$

**f** Using  $\cos(A+B) \equiv \cos A \cos B - \sin A \sin B$  gives  
 $\cos 4\theta \cos 3\theta - \sin 4\theta \sin 3\theta \equiv \cos(4\theta + 3\theta) \equiv \cos 7\theta$

**g** Using  $\sin(A+B) \equiv \sin A \cos B + \cos A \sin B$  gives  
 $\sin \frac{1}{2}\theta \cos 2\frac{1}{2}\theta + \cos \frac{1}{2}\theta \sin 2\frac{1}{2}\theta \equiv \sin\left(\frac{1}{2}\theta + 2\frac{1}{2}\theta\right) \equiv \sin 3\theta$

**h** Using  $\tan(A+B) \equiv \frac{\tan A + \tan B}{1 - \tan A \tan B}$  gives  
 $\frac{\tan 2\theta + \tan 3\theta}{1 - \tan 2\theta \tan 3\theta} \equiv \tan(2\theta + 3\theta) \equiv \tan 5\theta$

**i** Using  $\sin(P-Q) \equiv \sin P \cos Q - \cos P \sin Q$  gives  
 $\sin(A+B) \cos B - \cos(A+B) \sin B \equiv \sin((A+B) - B) \equiv \sin A$

**j** Using  $\cos(A+B) \equiv \cos A \cos B - \sin A \sin B$  gives  
 $\cos\left(\frac{3x+2y}{2}\right) \cos\left(\frac{3x-2y}{2}\right) - \sin\left(\frac{3x+2y}{2}\right) \sin\left(\frac{3x-2y}{2}\right) \equiv \cos\left(\left(\frac{3x+2y}{2}\right) + \left(\frac{3x-2y}{2}\right)\right)$   
 $\equiv \cos\left(\frac{6x}{2}\right) \equiv \cos 3x$

**10 a** Use the fact that  $\frac{1}{\sqrt{2}} = \cos \frac{\pi}{4} = \sin \frac{\pi}{4}$  to write

$$\frac{1}{\sqrt{2}}(\sin x + \cos x) = \frac{1}{\sqrt{2}}\sin x + \frac{1}{\sqrt{2}}\cos x = \sin x \cos \frac{\pi}{4} + \cos x \sin \frac{\pi}{4} = \sin\left(x + \frac{\pi}{4}\right)$$

or

$$\frac{1}{\sqrt{2}}(\sin x + \cos x) = \frac{1}{\sqrt{2}}\cos x + \frac{1}{\sqrt{2}}\sin x = \cos x \cos \frac{\pi}{4} + \sin x \sin \frac{\pi}{4} = \cos\left(x - \frac{\pi}{4}\right)$$

**b** Use the fact that  $\frac{1}{\sqrt{2}} = \cos \frac{\pi}{4} = \sin \frac{\pi}{4}$  to write

$$\frac{1}{\sqrt{2}}(\cos x - \sin x) = \frac{1}{\sqrt{2}}\cos x - \frac{1}{\sqrt{2}}\sin x = \cos x \cos \frac{\pi}{4} - \sin x \sin \frac{\pi}{4} = \cos\left(x + \frac{\pi}{4}\right)$$

**c** Use the fact that  $\frac{1}{2} = \cos \frac{\pi}{3} = \sin \frac{\pi}{6}$  and  $\frac{\sqrt{3}}{2} = \cos \frac{\pi}{6} = \sin \frac{\pi}{3}$  to write

$$\frac{1}{2}(\sin x + \sqrt{3}\cos x) = \frac{1}{2}\sin x + \frac{\sqrt{3}}{2}\cos x = \sin x \cos \frac{\pi}{3} + \cos x \sin \frac{\pi}{3} = \sin\left(x + \frac{\pi}{3}\right)$$

or

$$\frac{1}{2}(\sin x + \sqrt{3}\cos x) = \frac{\sqrt{3}}{2}\cos x + \frac{1}{2}\sin x = \cos x \cos \frac{\pi}{6} + \sin x \sin \frac{\pi}{6} = \cos\left(x - \frac{\pi}{6}\right)$$

**d** Use the fact that  $\frac{1}{\sqrt{2}} = \cos \frac{\pi}{4} = \sin \frac{\pi}{4}$  to write

$$\frac{1}{\sqrt{2}}(\sin x - \cos x) = \frac{1}{\sqrt{2}}\sin x - \frac{1}{\sqrt{2}}\cos x = \sin x \cos \frac{\pi}{4} - \cos x \sin \frac{\pi}{4} = \sin\left(x - \frac{\pi}{4}\right)$$

**11**  $\cos y = \sin(x + y)$

$$\Rightarrow \cos y = \sin x \cos y + \cos x \sin y$$

Divide throughout by  $\cos x \cos y$

$$\frac{\cancel{\cos y}^1}{\cos x \cancel{\cos y}} = \frac{\sin x \cancel{\cos y}}{\cos x \cancel{\cos y}} + \frac{\cancel{\cos x} \sin y}{\cancel{\cos x} \cos y}$$

$$\Rightarrow \sec x = \tan x + \tan y$$

$$\Rightarrow \tan y = \sec x - \tan x$$

**12** As  $\tan(x - y) = 3$

$$\text{so } \frac{\tan x - \tan y}{1 + \tan x \tan y} = 3$$

$$\Rightarrow \tan x - \tan y = 3 + 3 \tan x \tan y$$

$$\Rightarrow 3 \tan x \tan y + \tan y = \tan x - 3$$

$$\Rightarrow \tan y(3 \tan x + 1) = \tan x - 3$$

$$\Rightarrow \tan y = \frac{\tan x - 3}{3 \tan x + 1}$$

$$13 \sin x(\cos y + 2 \sin y) = \cos x(2 \cos y - \sin y)$$

$$\Rightarrow \sin x \cos y + 2 \sin x \sin y = 2 \cos x \cos y - \cos x \sin y$$

$$\Rightarrow \sin x \cos y + \cos x \sin y = 2(\cos x \cos y - \sin x \sin y)$$

$$\Rightarrow \sin(x + y) = 2 \cos(x + y)$$

$$\Rightarrow \frac{\sin(x + y)}{\cos(x + y)} = 2$$

$$\Rightarrow \tan(x + y) = 2$$

$$14 \text{ a } \tan(x - 45^\circ) = \frac{1}{4}$$

$$\Rightarrow \frac{\tan x - \tan 45^\circ}{1 + \tan x \tan 45^\circ} = \frac{1}{4}$$

$$\Rightarrow 4 \tan x - 4 = 1 + \tan x \quad (\text{as } \tan 45^\circ = 1)$$

$$\Rightarrow 3 \tan x = 5$$

$$\Rightarrow \tan x = \frac{5}{3}$$

$$\text{b } \sin(x - 60^\circ) = 3 \cos(x + 30^\circ)$$

$$\Rightarrow \sin x \cos 60^\circ - \cos x \sin 60^\circ = 3 \cos x \cos 30^\circ - 3 \sin x \sin 30^\circ$$

$$\Rightarrow \frac{1}{2} \sin x - \frac{\sqrt{3}}{2} \cos x = \frac{3\sqrt{3}}{2} \cos x - \frac{3}{2} \sin x$$

$$\Rightarrow 4 \sin x = 4\sqrt{3} \cos x$$

$$\Rightarrow \frac{\sin x}{\cos x} = \frac{4\sqrt{3}}{4}$$

$$\Rightarrow \tan x = \sqrt{3}$$

$$\text{c } \tan(x - 60^\circ) = 2$$

$$\Rightarrow \frac{\tan x - \tan 60^\circ}{1 + \tan x \tan 60^\circ} = 2$$

$$\Rightarrow \frac{\tan x - \sqrt{3}}{1 + \sqrt{3} \tan x} = 2 \quad (\text{as } \tan 60^\circ = \sqrt{3})$$

$$\Rightarrow \tan x - \sqrt{3} = 2 + 2\sqrt{3} \tan x$$

$$\Rightarrow (2\sqrt{3} - 1) \tan x = -(2 + \sqrt{3})$$

$$\begin{aligned} \Rightarrow \tan x &= -\frac{(2 + \sqrt{3})}{2\sqrt{3} - 1} = -\frac{(2 + \sqrt{3})(2\sqrt{3} + 1)}{(2\sqrt{3} - 1)(2\sqrt{3} + 1)} \\ &= -\frac{8 + 5\sqrt{3}}{11} \end{aligned}$$

$$\begin{aligned}
 \mathbf{15} \quad \tan\left(x + \frac{\pi}{3}\right) &= \frac{1}{2} \\
 \Rightarrow \frac{\tan x + \tan \frac{\pi}{3}}{1 - \tan x \tan \frac{\pi}{3}} &= \frac{1}{2} \\
 \Rightarrow \frac{\tan x + \sqrt{3}}{1 - \sqrt{3} \tan x} &= \frac{1}{2} \quad \left(\tan \frac{\pi}{3} = \sqrt{3}\right) \\
 \Rightarrow 2 \tan x + 2\sqrt{3} &= 1 - \sqrt{3} \tan x \\
 \Rightarrow (2 + \sqrt{3}) \tan x &= 1 - 2\sqrt{3} \\
 \Rightarrow \tan x &= \frac{1 - 2\sqrt{3}}{2 + \sqrt{3}} = \frac{(1 - 2\sqrt{3})(2 - \sqrt{3})}{(2 + \sqrt{3})(2 - \sqrt{3})} \\
 &= \frac{2 - 4\sqrt{3} - \sqrt{3} + 6}{1} = 8 - 5\sqrt{3}
 \end{aligned}$$

$$\mathbf{16} \quad \text{Write } \theta = \left(\theta + \frac{2\pi}{3}\right) - \frac{2\pi}{3} \text{ and } \theta + \frac{4\pi}{3} = \left(\theta + \frac{2\pi}{3}\right) + \frac{2\pi}{3}$$

Now use the appropriate addition formulae for cos

$$\cos\left(\left(\theta + \frac{2\pi}{3}\right) - \frac{2\pi}{3}\right) = \cos\left(\theta + \frac{2\pi}{3}\right) \cos \frac{2\pi}{3} + \sin\left(\theta + \frac{2\pi}{3}\right) \sin \frac{2\pi}{3}$$

$$\cos\left(\left(\theta + \frac{2\pi}{3}\right) + \frac{2\pi}{3}\right) = \cos\left(\theta + \frac{2\pi}{3}\right) \cos \frac{2\pi}{3} - \sin\left(\theta + \frac{2\pi}{3}\right) \sin \frac{2\pi}{3}$$

Now add up all terms

$$\begin{aligned}
 \cos \theta + \cos\left(\theta + \frac{2\pi}{3}\right) + \cos\left(\theta + \frac{4\pi}{3}\right) \\
 &\equiv \cos\left(\left(\theta + \frac{2\pi}{3}\right) - \frac{2\pi}{3}\right) + \cos\left(\theta + \frac{2\pi}{3}\right) + \cos\left(\left(\theta + \frac{2\pi}{3}\right) + \frac{2\pi}{3}\right) \\
 &\equiv \cos\left(\theta + \frac{2\pi}{3}\right) \cos \frac{2\pi}{3} + \sin\left(\theta + \frac{2\pi}{3}\right) \sin \frac{2\pi}{3} + \cos\left(\theta + \frac{2\pi}{3}\right) \cos \frac{2\pi}{3} \\
 &\quad - \sin\left(\theta + \frac{2\pi}{3}\right) \sin \frac{2\pi}{3} \\
 &\equiv 2 \cos\left(\theta + \frac{2\pi}{3}\right) \cos \frac{2\pi}{3} + \cos\left(\theta + \frac{2\pi}{3}\right) \\
 &\equiv 0 \text{ as } \cos \frac{2\pi}{3} = -\frac{1}{2}
 \end{aligned}$$

**Challenge**

$$\begin{aligned}\mathbf{a \ i} \quad \text{Area} &= \frac{1}{2} ab \sin \theta = \frac{1}{2} x(y \cos B)(\sin A) \\ &= \frac{1}{2} xy \sin A \cos B\end{aligned}$$

$$\begin{aligned}\mathbf{ii} \quad \text{Area} &= \frac{1}{2} ab \sin \theta = \frac{1}{2} y(x \cos A)(\sin B) \\ &= \frac{1}{2} xy \cos A \sin B\end{aligned}$$

$$\mathbf{iii} \quad \text{Area} = \frac{1}{2} ab \sin \theta = \frac{1}{2} xy \sin(A + B)$$

$$\begin{aligned}\mathbf{b} \quad \text{Area } T_1 + T_2 &= \text{Area } T_1 + \text{Area } T_2 \\ \Rightarrow \frac{1}{2} xy \sin(A + B) &= \frac{1}{2} xy \sin A \cos B + \frac{1}{2} xy \cos A \sin B \\ \Rightarrow \sin(A + B) &= \sin A \cos B + \cos A \sin B\end{aligned}$$