

Dynamics 5C

1 a $v^2 = \omega^2 (a^2 - x^2)$

$a = 0.5, \quad x = 0 \quad c = 2$

$2^2 = \omega^2 \times 0.5^2$

$\omega = \frac{2}{0.5} = 4$

period = $\frac{2\pi}{\omega} = \frac{2\pi}{4} = \frac{\pi}{2}$

The period is $\frac{\pi}{2}$ s.

b $x = 0.2 \text{ m} \quad v^2 = 4^2 (0.5^2 - 0.2^2)$

$v = 1.833\dots$

When $OP = 0.2 \text{ m}$ the speed of P is 1.83 m s^{-1} (3 s.f.)

2 a period = $\frac{2\pi}{\omega} = \frac{\pi}{3}$

$\therefore \omega = 6$

$v^2 = \omega^2 (a^2 - x^2)$

$6^2 = 6^2 (a^2 - 0^2)$

$\therefore a = 1$

The amplitude is 1 m.

Maximum speed occurs when $x = 0$.

b $x = a \sin \omega t$

$v = a\omega \cos \omega t$

$t = 0.3 \text{ s} \quad v = 1 \times 6 \cos(6 \times 0.3)$

$v = 6 \cos 1.8$

$v = -1.363$

The speed 0.3 s after passing O is required.

Differentiate the line above to obtain v .

when $t = 0.3$, P has speed 1.36 m s^{-1} (3 s.f.)

Speed is positive.

3 a $v^2 = \omega^2 (a^2 - x^2)$

$x = 0, \quad v = 10 \text{ m s}^{-1} \quad 10^2 = \omega^2 a^2 \quad (1)$

$\ddot{x} = -\omega^2 x$

$x = -a, \quad \ddot{x} = 20 \text{ m s}^{-2}$

$20 = +\omega^2 a$

(2)

(1) \div (2) $\frac{100}{20} = \frac{\omega^2 a^2}{\omega^2 a}$

$a = 5$

Maximum acceleration occurs when $x = -a$.

The amplitude is 5 m.

3 b Using (1) $10 = a\omega$

$$10 = 5\omega$$

$$\omega = 2$$

$$\text{period} = \frac{2\pi}{\omega} = \pi$$

The period is π s.

4 $\text{period} = \frac{2\pi}{\omega} = \frac{3\pi}{5}$

$$\omega = \frac{10}{3}$$

$$v^2 = \omega^2(a^2 - x^2)$$

$$v^2 = \left(\frac{10}{3}\right)^2 (0.4^2 - 0)$$

$$v = \frac{10}{3} \times 0.4 = \frac{4}{3}$$

The maximum speed is $\frac{4}{3} \text{ m s}^{-1}$.

Maximum speed occurs when $x = 0$.

5 $\ddot{x} = -\omega^2 x$

$$\ddot{x} = 15 \text{ m s}^{-2}, x = a$$

$$15 = \omega^2 a \quad (1)$$

$$v^2 = \omega^2(a^2 - x^2)$$

$$v = 18 \text{ m s}^{-1}, x = 0 \quad 18^2 = \omega^2 a^2 \quad (2)$$

$$(2) \div (1) \quad \frac{18^2}{15} = \frac{\omega^2 a^2}{\omega^2 a}$$

$$a = \frac{18^2}{15} = 21.6$$

Using (2) $a\omega = 18$

$$\omega = \frac{18}{21.6} = 0.8333\dots$$

$$v^2 = \omega^2(a^2 - x^2)$$

$$v^2 = 0.8333\dots^2 (21.6^2 - 2.5^2)$$

$$v = 17.87\dots$$

The speed is 17.9 m s^{-1} (3 s.f.)

First find a and ω (see question 3.)

6 a $\text{period} = \frac{2\pi}{\omega} = \frac{\pi}{2}$
 $\omega = 4$

$$v^2 = \omega^2(a^2 - x^2)$$

$$x = 1.2 \text{ m} \quad v = 1.5 \text{ ms}^{-1}$$

$$1.5^2 = 4^2(a^2 - 1.2^2)$$

$$a^2 = \frac{1.5^2}{4^2} + 1.2^2$$

$$a = 1.257\dots$$

The amplitude is 1.26 m (3 s.f.).



Use the period to find ω .



Then use $v^2 = \omega^2(a^2 - x^2)$ with $x = 1.2$ and $v = 1.5$ to find a .

b $x = a \sin \omega t$

$$x = 1.26 \sin 4t$$



Use $x = a \sin \omega t$ as $x = 0$ when $t = 0$.

7 a $\text{period} = \frac{2\pi}{\omega} = \frac{1}{6}$

$$\omega = 12\pi$$

$$v^2 = \omega^2(a^2 - x^2)$$

$$5^2 = (12\pi)^2(a^2 - 0)$$

$$a = \frac{5}{12\pi} = 0.1326\dots$$

The amplitude is 0.133 m (3 s.f.)



The period is the time for one complete oscillation.



You are told the magnitude of the acceleration at A .

b $\ddot{x} = -\omega^2 x$

$$20 = |-12^2 \pi^2| x$$

$$x = \frac{20}{144\pi^2}$$

$$x = 0.01407\dots$$

$$OA = 0.0141 \text{ m (3 s.f.)}$$

8 a $v^2 = \omega^2(a^2 - x^2)$
 $x = 0.6 \text{ m}, v = 3 \text{ m s}^{-1}$
 $3^2 = \omega^2(a^2 - 0.6^2)$ (1)

$x = 0.2 \text{ m}, v = 6 \text{ m s}^{-1}$
 $6^2 = \omega^2(a^2 - 0.2^2)$ (2)

(2) \div (1) $\frac{6^2}{3^2} = \frac{\omega^2(a^2 - 0.2^2)}{\omega^2(a^2 - 0.6^2)}$

$4(a^2 - 0.6^2) = a^2 - 0.2^2$
 $3a^2 = 4 \times 0.6^2 - 0.2^2$
 $a^2 = \frac{4 \times 0.6^2 - 0.2^2}{3}$
 $a = 0.6831\dots$

The distance AB is 1.37 m (3 s.f.)

← AB is twice the amplitude.

b Using (1) $9 = \omega^2(0.6831^2 - 0.6^2)$

$\omega^2 = \frac{9}{(0.6831^2 - 0.6^2)}$

$\omega = 9.187$

period = $\frac{2\pi}{\omega} = \frac{2\pi}{9.187} = 0.6838\dots$

The period is 0.684s (3 s.f.).

9 a period = $\frac{2\pi}{\omega} = 2\pi$

$\omega = 1$

$v^2 = \omega^2(a^2 - x^2)$

$x = 1 \text{ m}, v = 0.1 \text{ m s}^{-1}$

$0.1^2 = 1^2(a^2 - 1^2)$

$a^2 = 0.1^2 + 1^2$

$a = 1.004\dots$

$v_{\max} = \omega a$

$= 1 \times 1.004\dots$

The maximum speed is 1.00 m s⁻¹ (3 s.f.).

← First find ω .

← Now use $v^2 = \omega^2(a^2 - x^2)$ to find a .

← Maximum speed occurs when $x = 0$.

b $v^2 = 1(1.004^2 - 0.4^2)$

$v = 0.9219\dots$

The speed is 0.922 m s⁻¹ (3 s.f.).

$$10 \quad a = \frac{2.5}{2} = 1.25$$

$$\text{Period} = \frac{2\pi}{\omega} = \frac{60}{30} = 2$$

$$\omega = \pi$$

$$v_{\max} = a\omega$$

$$= 1.25 \times \pi$$

$$\text{maximum K.E.} = \frac{1}{2}mv_{\max}^2$$

$$= \frac{1}{2} \times 1.2 \times 1.25^2 \times \pi^2$$

$$= 9.252\dots$$

The maximum K.E. is 9.25J (3s.f.).

30 oscillations per minute \Rightarrow
2s for 1 oscillation

$$11 \text{ a} \quad a = 0.8 \div 2 = 0.4 \text{ m}$$

$$\text{period} = \frac{2\pi}{\omega} = 2$$

$$\omega = \pi$$

$$v^2 = \omega^2 (a^2 - x^2)$$

$$x = 0 \quad v = \omega a$$

$$v = \pi \times 0.4$$

$$v = 1.256\dots$$

The maximum speed is 1.26 m s⁻¹ (3 s.f.).

The amplitude is half the distance
between the highest and lowest points.

b 0.6 m from highest point

$$\Rightarrow x = -0.2 \text{ m}$$

$$x = a \cos \omega t$$

$$-0.2 = 0.4 \cos \pi t$$

$$\cos \pi t = -0.5$$

$$t = \frac{1}{\pi} \cos^{-1}(-0.5)$$


$$t = \frac{1}{\pi} \times \left(\pi - \frac{\pi}{3} \right)$$

$$t = \frac{2}{3}$$

The buoy takes $\frac{2}{3}$ s to fall 0.6 m.

The buoy is now below the centre.

You want the time from the highest point.



$$\text{period} = \frac{2\pi}{\omega} = 2$$

$$\therefore \omega = \pi$$

$$x = a \sin \omega t$$

$$x = 0.5 \sin \pi t$$

$$x = 0.2 \text{ m } 0.2 = 0.5 \sin \pi t_1$$

$$\pi t_1 = \sin^{-1}\left(\frac{0.2}{0.5}\right) = \sin^{-1}\left(\frac{2}{5}\right)$$

$$x = 0.3 \quad \pi t_2 = \sin^{-1}\left(\frac{3}{5}\right)$$

$$\text{time } A \rightarrow B = t_2 - t_1$$

$$= \frac{1}{\pi} \left(\sin^{-1}\left(\frac{3}{5}\right) - \sin^{-1}\left(\frac{2}{5}\right) \right)$$

$$= 0.07384\dots$$

The time to move directly from A to B is 0.0738 s (3 s.f.).

Use $x = a \sin \omega t$ to find the time to go from O to A and the time to go from O to B .

13 a $x = 4 \sin 2t$

$$\dot{x} = 8 \cos 2t$$

$$\ddot{x} = -16 \sin 2t$$

$$\ddot{x} = -4(4 \sin 2t)$$

$$\ddot{x} = -4x$$

\therefore S.H.M.

Differentiate the given equation twice.

b amplitude = 4 m

$$\text{period} = \frac{2\pi}{2} = \pi \text{ s}$$

Compare $x = 4 \sin 2t$ with $x = a \sin \omega t$ to obtain a and ω .

c $v^2 = \omega^2(a^2 - x^2)$

$$x = 0 \quad v^2 = 4(4^2 - 0)$$

$$v = 8$$

The maximum speed is 8 m s^{-1} .

d $x = 4 \sin 2t$

$$\dot{x} = 8 \cos 2t$$

$$\dot{x} = 4 \text{ ms}^{-1} \quad 4 = 8 \cos 2t$$

$$\cos 2t = 0.5$$

$$t = \frac{1}{2} \cos^{-1} 0.5$$

$$t = \frac{1}{2} \times \frac{\pi}{3}$$

The least value of t is $\frac{\pi}{6}$.

From a.

$$13 \text{ e} \quad x = 4 \sin 2t$$

$$x = 2 \quad 2 = 4 \sin 2t$$

$$\sin 2t = 0.5$$

$$t = \frac{1}{2} \sin^{-1} 0.5$$

$$t = \frac{1}{2} \times \frac{\pi}{6}$$

The least value of t is $\frac{\pi}{12}$.

$$14 \text{ a} \quad x = 3 \sin\left(4t + \frac{1}{2}\right)$$

$$\dot{x} = 12 \cos\left(4t + \frac{1}{2}\right)$$

$$\ddot{x} = -48 \sin\left(4t + \frac{1}{2}\right)$$

$$\ddot{x} = -16x$$

\therefore S.H.M.

$$\text{b} \quad \text{amplitude} = 3 \text{ m}$$

$$\text{period} = \frac{2\pi}{\omega} = \frac{2\pi}{4} = \frac{\pi}{2} \text{ s}$$

Compare with $x = a \sin(\omega t + \varepsilon)$
to obtain a and ω .

$$\text{c} \quad t = 0 \quad x = 3 \sin\left(\frac{1}{2}\right)$$

$$= 1.438\dots$$

When $t = 0$, $x = 1.44$ (3 s.f.)

$$\text{d} \quad x = 0 \quad 0 = 3 \sin\left(4t + \frac{1}{2}\right)$$

$$\sin\left(4t + \frac{1}{2}\right) = 0$$

$$4t + \frac{1}{2} = 0, \pi, \dots$$

$$4t = \left(0 - \frac{1}{2}\right), \left(\pi - \frac{1}{2}\right), \dots$$

$$t = -\frac{1}{8} \text{ (not applicable)}$$

$$t = \frac{1}{4} \left(\pi - \frac{1}{2}\right) = 0.6603\dots$$

The value of t is 0.660 (3 s.f.).

15 a amplitude = $\frac{(15-5)}{2} = 5 \text{ m}$

The difference between high and low tides is twice the amplitude.

period = $(10 \text{ am} \rightarrow 4.15 \text{ pm}) \times 2$
 $= 6.25 \times 2 \text{ hr}$
 $= 12.5 \text{ hr}$

The time from low to high tide is half the period.

You can work through this question using hours as the unit for time.


period = $\frac{2\pi}{\omega} = 12.5$

$\omega = \frac{2\pi}{12.5}$

$x = a \cos \omega t$

Start at low tide.

$x = 5 \cos\left(\frac{2\pi}{12.5}t\right)$



$x = 3 \text{ m}$

$3 = 5 \cos\left(\frac{2\pi}{12.5}t\right)$

The diagram shows that when the water is 7 m deep, $x = 3$.

$\cos\left(\frac{2\pi}{12.5}t\right) = 0.6$

$t = \frac{12.5}{2\pi} \cos^{-1} 0.6$

$t = 1.844\dots$

Time after 10 am.

The ship can enter the harbour at 11.51 am. (nearest minute).

b The water is once more 7 m deep at $(12.5 - 1.844)$ hours after 10 am

$= 10.656 \text{ hrs after 10 am}$

$= 10 \text{ hr } 39.3\dots \text{min.}$

\therefore Ship must leave by 8.39 pm (nearest minute).

Remember to change the decimal part of 1.844 into minutes (0.844×60) .

Use the symmetry of S.H.M. to find the time required.

16 $\overline{A \quad 0.4 \text{ m} \quad O \quad 0.5 \text{ m} \quad B}$

$$\text{period} = \frac{2\pi}{\omega} = 4$$

$$\omega = \frac{\pi}{2}$$

$$x = a \sin \omega t$$

$$x = 0.75 \sin \frac{\pi}{2} t$$

$$x = 0.5 \text{ m} \quad 0.5 = 0.75 \sin \frac{\pi}{2} t$$

$$\sin \frac{\pi t}{2} = \frac{0.5}{0.75}$$

$$t = \frac{2}{\pi} \sin^{-1} \left(\frac{0.5}{0.75} \right)$$

$$x = 0.4 \text{ m} \quad t = \frac{2}{\pi} \sin^{-1} \left(\frac{0.4}{0.75} \right)$$

Time $B \rightarrow A$

$$= \frac{2}{\pi} \left[\sin^{-1} \left(\frac{0.5}{0.75} \right) + \sin^{-1} \left(\frac{0.4}{0.75} \right) \right]$$

$$= 0.8226\dots$$

P takes 0.823s to travel directly from B to A (3 s.f.)

Find the time taken from O to B (using $x = 0.5$ m) and from O to the point where $x = 0.4$ m.

Adding these times will give the time to go directly from B to A due to the symmetry of S.H.M.

Challenge

$$\ddot{x} = -\omega^2 x \qquad v^2 = \omega^2 (a^2 - x^2)$$

$$v_1^2 = \omega^2 (a^2 - x_1^2) \quad (1)$$

$$v_2^2 = \omega^2 (a^2 - x_2^2) \quad (2)$$

$$(2) - (1): v_2^2 - v_1^2 = \omega^2 (a^2 - x_2^2) - \omega^2 (a^2 - x_1^2)$$

$$v_2^2 - v_1^2 = \omega^2 (a^2 - x_2^2 - a^2 + x_1^2)$$

$$\text{Rearranging gives } \omega^2 = \frac{v_2^2 - v_1^2}{x_1^2 - x_2^2} \text{ so } \omega^2 = \left(\frac{v_2^2 - v_1^2}{x_1^2 - x_2^2} \right)^{\frac{1}{2}}$$

$$T = \frac{2\pi}{\omega} = 2\pi \left(\frac{x_1^2 - x_2^2}{v_2^2 - v_1^2} \right)^{\frac{1}{2}}$$