

## Chapter review 4

1 a  $y = x^2(x - 2)$

$$0 = x^2(x - 2)$$

$$\text{So } x = 0 \text{ or } x = 2$$

The curve crosses the  $x$ -axis at  $(2, 0)$

and touches it at  $(0, 0)$ .

$$x \rightarrow \infty, y \rightarrow \infty$$

$$x \rightarrow -\infty, y \rightarrow -\infty$$

$$y = 2x - x^2$$

$$= x(2 - x)$$


As  $a = -1$  is negative, the graph has a  $\wedge$  shape and a maximum point.

$$0 = x(2 - x)$$

$$\text{So } x = 0 \text{ or } x = 2$$

The curve crosses the  $x$ -axis at  $(0, 0)$

and  $(2, 0)$ .



b  $x^2(x - 2) = x(2 - x)$

$$x^2(x - 2) - x(2 - x) = 0$$

$$x^2(x - 2) + x(x - 2) = 0$$

$$x(x - 2)(x + 1) = 0$$

$$\text{So } x = 0, x = 2 \text{ or } x = -1$$

Using  $y = x(2 - x)$ :

$$\text{when } x = 0, y = 0 \times 2 = 0$$

$$\text{when } x = 2, y = 2 \times 0 = 0$$


$$\text{when } x = -1, y = (-1) \times 3 = -3$$

The points of intersection are  $(0, 0)$ ,

$(2, 0)$  and  $(-1, -3)$ .

2 a  $y = \frac{6}{x}$  is like  $y = \frac{1}{x}$ .

$y = 1 + x$  is a straight line.



2 b  $\frac{6}{x} = 1 + x$

$$6 = x + x^2$$

$$0 = x^2 + x - 6$$

$$0 = (x + 3)(x - 2)$$

$$\text{So } x = 2 \text{ or } x = -3$$

Using  $y = 1 + x$ :

$$\text{when } x = 2, y = 1 + 2 = 3$$

$$\text{when } x = -3, y = 1 - 3 = -2$$

So  $A$  is  $(-3, -2)$  and  $B$  is  $(2, 3)$ .

c Substituting the points  $A$  and  $B$  into

$$y = x^2 + px + q:$$

$$A: -2 = 9 - 3p + q \quad (1)$$

$$B: 3 = 4 + 2p + q \quad (2)$$

$$(1) - (2):$$

$$-5 = 5 - 5p$$

$$p = 2$$

Substituting in (1):

$$-2 = 9 - 6 + q$$

$$q = -5$$


d  $y = x^2 + 2x - 5$

As  $a = 1$  is positive, the graph has a  $\vee$  shape and a minimum point.

$$y = (x + 1)^2 - 6$$

So the minimum is at  $(-1, -6)$ .

3 a  $f(2x)$  is a stretch with scale factor  $\frac{1}{2}$  in the  $x$ -direction.




$$A'(\frac{3}{2}, 4), B'(0, 0)$$

The asymptote is  $y = 2$ .


b  $\frac{1}{2}f(x)$  is a stretch with scale factor  $\frac{1}{2}$  in the  $y$ -direction.

3 b




$A'(3, 2), B'(0, 0)$   
The asymptote is  $y = 1$ .

c  $f(x) - 2$  is a translation by  $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$ ,  
or two units down.




$A'(3, 2), B'(0, -2)$   
The asymptote is  $y = 0$ .

d  $f(x + 3)$  is a translation by  $\begin{pmatrix} -3 \\ 0 \end{pmatrix}$ ,  
or three units to the left.




$A'(0, 4), B'(-3, 0)$   
The asymptote is  $y = 2$ .

e  $f(x - 3)$  is a translation by  $\begin{pmatrix} 3 \\ 0 \end{pmatrix}$ ,  
or three units to the right.



e  $A'(6, 4), B'(3, 0)$   
The asymptote is  $y = 2$ .


f  $f(x) + 1$  is a translation by  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ,  
or one unit up.




$A'(3, 5), B'(0, 1)$   
The asymptote is  $y = 3$ .

4  $2 = 5 + 2x - x^2$   
 $x^2 - 2x - 3 = 0$   
 $(x - 3)(x + 1) = 0$   
So  $x = -1$  or  $x = 3$   
The points of intersection  
are  $A(-1, 2), B(3, 2)$ .


5 a  $f(-x)$  is a reflection in the  $y$ -axis.




b  $-f(x)$  is a reflection in the  $x$ -axis.



- 6 a** Let  $y = a(x - p)(x - q)$ .  
 Since  $(1, 0)$  and  $(3, 0)$  are on the curve then  
 $p = 1$  and  $q = 3$ .  
 So  $y = a(x - 1)(x - 3)$   
 Using  $(2, -1)$ :  
 $-1 = a(1)(-1)$   
 $a = 1$   
 So  $y = (x - 1)(x - 3) = x^2 - 4x + 3$
- b i**  $f(x + 2) = (x + 1)(x - 1)$ , or a translation  
 by  $\begin{pmatrix} -2 \\ 0 \end{pmatrix}$ , or two units to the left.



- ii**  $f(2x) = (2x - 1)(2x - 3)$ , or a stretch with  
 scale factor  $\frac{1}{2}$  in the  $x$ -direction.



- 7 a**  $f(x) = (x - 1)(x - 2)(x + 1)$   
 When  $x = 0$ ,  $y = (-1) \times (-2) \times 1 = 2$   
 So the curve crosses the  $y$ -axis at  $(0, 2)$ .
- b**  $y = af(x)$  is a stretch with scale factor  $a$   
 in the  $y$ -direction.  
 The  $y$ -coordinate has multiplied by  $-2$ ,  
 therefore  $y = -2f(x)$ .  
 $a = -2$

- 7 c**  $f(x) = (x - 1)(x - 2)(x + 1)$   
 $0 = (x - 1)(x - 2)(x + 1)$   
 So  $x = 1$ ,  $x = 2$  or  $x = -1$   
 The curve crosses the  $x$ -axis at  $(1, 0)$ ,  
 $(2, 0)$  and  $(-1, 0)$ .  
 $y = f(x + b)$  is a translation  $b$  units  
 to the left.  
 For the point  $(0, 0)$  to lie on the translated  
 curve, either the point  $(1, 0)$ ,  $(2, 0)$  or  
 $(-1, 0)$  has translated to the point  $(1, 0)$ .  
 For the coordinate  $(1, 0)$  to be translated to  
 $(0, 0)$ ,  $b = 1$ .  
 For the coordinate  $(2, 0)$  to be translated to  
 $(0, 0)$ ,  $b = 2$ .  
 For the coordinate  $(-1, 0)$  to be translated to  
 $(0, 0)$ ,  $b = -1$ .  
 $b = -1$ ,  $b = 1$  or  $b = 2$

- 8 a i**  $y = f(3x)$  is a stretch with scale factor  $\frac{1}{3}$   
 in the  $x$ -direction. Find  $\frac{1}{3}$  of the  
 $x$ -coordinate.  
 $P$  is transformed to  $(\frac{4}{3}, 3)$ .

- ii**  $\frac{1}{2}y = f(x)$   
 $y = 2f(x)$ , which is a stretch with scale  
 factor 2 in the  $y$ -direction.  
 $P$  is transformed to  $(4, 6)$ .

- iii**  $y = f(x - 5)$  is a translation by  $\begin{pmatrix} 5 \\ 0 \end{pmatrix}$ ,  
 or five units to the right.  
 $P$  is transformed to  $(9, 3)$ .

- iv**  $-y = f(x)$   
 $y = -f(x)$ , which is a reflection of the  
 curve in the  $x$ -axis.  
 $(4, -3)$

- v**  $2(y + 2) = f(x)$   
 $y = \frac{1}{2}f(x) - 2$ , which is a stretch with  
 scale factor  $\frac{1}{2}$  in the  $y$ -direction and  
 then a translation by  $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$ , or two  
 units down.  
 $P$  is transformed to  $(4, -\frac{1}{2})$ .


- 8 b**  $P(4, 3)$  is transformed to  $(2, 3)$ .  
 Either the  $x$ -coordinate has halved, which is a stretch with scale factor  $\frac{1}{2}$  in the  $x$ -direction, or it has had 2 subtracted from it, which is a translation by  $\begin{pmatrix} -2 \\ 0 \end{pmatrix}$ , or two units to the left.

So the transformation is  $y = f(2x)$  or  $y = f(x + 2)$ .

- c i**  $P(4, 3)$  is translated to the point  $(8, 6)$ .  
 The  $x$ -coordinate of  $P$  has 4 added to it and the  $y$ -coordinate has 3 added to it.  
 $y = f(x - 4) + 3$
- ii**  $P(4, 3)$  is stretched to the point  $(8, 6)$ .  
 The  $x$ -coordinate of  $P$  has doubled and the  $y$ -coordinate has doubled.  
 $y = 2f(\frac{1}{2}x)$

**9 a**  $x^3 - 6x^2 + 9x$   
 $= x(x^2 - 6x + 9)$   
 $= x(x - 3)^2$

- b**  $y = x(x - 3)^2$   
 $0 = x(x - 3)^2$   
 So  $x = 0$  or  $x = 3$   
 The curve crosses the  $x$ -axis at  $(0, 0)$  and touches it at  $(3, 0)$ .  
 $x \rightarrow \infty, y \rightarrow \infty$   
 $x \rightarrow -\infty, y \rightarrow -\infty$



- c**  $y = (x - k)^3 - 6(x - k)^2 + 9(x - k)$  is a translation of the curve  $y = x^3 - 6x^2 + 9x$  by  $\begin{pmatrix} k \\ 0 \end{pmatrix}$ , or  $k$  units to the right.


For the point  $(-4, 0)$  to lie on the translated curve, either the point  $(0, 0)$  or  $(3, 0)$  has translated to the point  $(-4, 0)$ .

For the coordinate  $(0, 0)$  to be translated to  $(-4, 0)$ ,  $k = -4$ .


For the coordinate  $(3, 0)$  to be translated to  $(-4, 0)$ ,  $k = -7$ .

$$k = -4 \text{ or } k = -7$$


- 10 a**  $y = x(x - 2)^2$   
 $0 = x(x - 2)^2$   
 So  $x = 0$  or  $x = 2$   
 The curve crosses the  $x$ -axis at  $(0, 0)$  and touches it at  $(2, 0)$ .  
 $x \rightarrow \infty, y \rightarrow \infty$   
 $x \rightarrow -\infty, y \rightarrow -\infty$



- b**  $y = f(x + 3)$  is a translation by vector  $\begin{pmatrix} -3 \\ 0 \end{pmatrix}$  of  $y = f(x)$ , or three units to the left.  
 So the curve crosses the  $x$ -axis at  $(-3, 0)$  and touches it at  $(-1, 0)$ .  
 When  $x = 3$ ,  $f(x) = 3(3 - 2)^2 = 3$   
 So  $f(x + 3)$  crosses the  $y$ -axis at  $(0, 3)$ .



- 11 a**  $y = f(x) - 2$  is a translation by  $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$ , or two units down.




The horizontal asymptote is  $y = -2$ .  
The vertical asymptote is  $x = 0$ .

- b** From the sketch, the curve crosses the  $x$ -axis.

$$\begin{aligned} y &= f(x) - 2 \\ &= \frac{1}{x} - 2 \\ 0 &= \frac{1}{x} - 2 \\ x &= \frac{1}{2} \end{aligned}$$

So the curve cuts the  $x$ -axis at  $(\frac{1}{2}, 0)$ .

- c**  $y = f(x + 3)$  is a translation by  $\begin{pmatrix} -3 \\ 0 \end{pmatrix}$ , or three units to the left.



- d** The horizontal asymptote is  $y = 0$ .  
The vertical asymptote is  $x = -3$ .

$$\begin{aligned} y &= f(x + 3) \\ &= \frac{1}{x + 3} \end{aligned}$$

When  $x = 0$ ,  $y = \frac{1}{3}$

So the curve cuts the  $y$ -axis at  $(0, \frac{1}{3})$ .

### Challenge

$$R(6, -4)$$

$$y = f(x + c) - d \text{ is a translation by } \begin{pmatrix} -c \\ 0 \end{pmatrix},$$

or  $c$  units to the left and a translation by  $\begin{pmatrix} 0 \\ -d \end{pmatrix}$ , or  $d$  units down.

So  $R$  is transformed to  $(6 - c, -4 - d)$ .