

### Exercise 4B

#### 1 Method 1

If  $x < 0$ ,  $F(x) = 0$  so  $F(0) = 0$

If  $0 \leq x \leq 2$

$$F(x) = F(0) + \int_0^x \frac{3t^2}{8} dt = \left[ \frac{3t^3}{24} \right]_0^x = \frac{3x^3}{24} - 0 = \frac{x^3}{8}$$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{3x^3}{8} & 0 \leq x \leq 2 \\ 1 & x > 2 \end{cases}$$

#### Method 2

If  $0 \leq x \leq 2$

$$F(x) = \int \frac{3x^2}{8} dx = \frac{x^3}{8} + c$$

$$F(2) = 1 \Rightarrow 1 + c = 1 \Rightarrow c = 0$$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{3x^3}{8} & 0 \leq x \leq 2 \\ 1 & x > 2 \end{cases}$$

#### 2 Method 1

If  $x < 1$ ,  $F(x) = 0$  so  $F(1) = 0$

If  $1 \leq x \leq 3$

$$F(x) = F(1) + \int_1^x \frac{1}{4}(4-t) dt = \left[ t - \frac{t^2}{8} \right]_1^x = \left( x - \frac{x^2}{8} \right) - \left( 1 - \frac{1}{8} \right) = x - \frac{x^2}{8} - \frac{7}{8}$$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 1 \\ x - \frac{x^2}{8} - \frac{7}{8} & 1 \leq x \leq 3 \\ 1 & x > 3 \end{cases}$$

#### Method 2

If  $1 \leq x \leq 3$

$$F(x) = \int \frac{1}{4}(4-x) dx = x - \frac{x^2}{8} + c$$

$$F(3) = 1 \Rightarrow 3 - \frac{9}{8} + c = 1 \Rightarrow c = -\frac{7}{8}$$

So  $F(x) = x - \frac{x^2}{8} - \frac{7}{8}$ , which leads to the full solution given for Method 1.

3 If  $x \leq 0$ ,  $F(x) = 0$  so  $F(0) = 0$

If  $0 < x < 3$

$$F(x) = \int \frac{1}{9} x dx = \frac{1}{18} x^2 + c$$

$$\text{As } F(0) = 0 \Rightarrow c = 0$$

If  $3 \leq x \leq 6$

$$F(x) = \int \frac{1}{9} (6-x) dx = \frac{2x}{3} - \frac{x^2}{18} + d$$


$$\text{As } F(6) = 1 \Rightarrow 4 - 2 + d = 1 \Rightarrow d = -1$$

So the full solution is:

$$F(x) = \begin{cases} 0 & x \leq 0 \\ \frac{x^2}{18} & 0 < x < 3 \\ \frac{2x}{3} - \frac{x^2}{18} - 1 & 3 \leq x \leq 6 \\ 1 & x > 6 \end{cases}$$

This shows the solution using Method 2; the problem can also be solved using Method 1.

4 a The graph is a horizontal line from  $(0, k)$  to  $(3, k)$ , and a straight line from  $(3, k)$  to  $(5, 5k)$ . Otherwise  $f(x)$  is 0.



b The area under the curve must equal 1, so:

$$\int_0^3 k dx + \int_3^5 k(2x-5) dx = 1$$

$$k[x]_0^3 + k[(x^2-5x)]_3^5 = 1$$

$$3k + k((25-25) - (9-15)) = 1$$

$$9k = 1$$

$$k = \frac{1}{9}$$

4 c If  $x < 0$ ,  $F(x) = 0$  so  $F(0) = 0$

If  $0 \leq x < 3$

$$F(x) = \int \frac{1}{9} dx = \frac{x}{9} + c$$

As  $F(0) = 0 \Rightarrow c = 0$

If  $3 \leq x \leq 5$

$$F(x) = \int \frac{1}{9}(2x - 5) dx = \frac{x^2}{9} - \frac{5x}{9} + d$$

As  $F(5) = 1 \Rightarrow \frac{25}{9} - \frac{25}{9} + d = 1 \Rightarrow d = 1$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{x}{9} & 0 \leq x < 3 \\ \frac{x^2}{9} - \frac{5x}{9} + 1 & 3 \leq x \leq 5 \\ 1 & x > 5 \end{cases}$$

5  $f(x) = \frac{d}{dx} F(x)$

So where  $F(x)$  is constant,  $f(x) = 0$

For  $2 \leq x \leq 3$ ,  $f(x) = \frac{d}{dx} \frac{1}{5}(x^2 - 4) = \frac{2x}{5}$

So the probability density function is:

$$f(x) = \begin{cases} \frac{2x}{5} & 2 \leq x \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

6 a  $P(X \leq 2.5) = F(2.5) = \frac{1}{2}(2.5 - 1) = 0.75$

b  $P(X > 1.5) = 1 - F(1.5) = 1 - \frac{1}{2}(1.5 - 1) = 0.75$

c  $P(1.5 \leq X \leq 2.5) = F(2.5) - F(1.5) = 0.75 - 0.25 = 0.5$

7 As  $F(x)$  is a cumulative distribution function,  $F(2) = 0$  and  $F(4) = 1$

$$F(2) = 0 \Rightarrow \frac{2^p}{6} + q = 0 \quad (1)$$

$$F(4) = 0 \Rightarrow \frac{4^p}{6} + q = 1 \quad (2)$$

Subtracting equation (1) from equation (2) gives:

$$\frac{4^p}{6} - \frac{2^p}{6} = 1 \Rightarrow 4^p - 2^p - 6 = 0$$

Let  $y = 2^p$ , then  $y^2 = 2^p \cdot 2^p = 4^p$  and the equation can be written as:

$$y^2 - y - 6 = 0 \Rightarrow (y - 3)(y + 2) = 0$$

Taking the positive root,  $y = 3 \Rightarrow 2^p = 3$

So taking logs of both sides,  $\ln 2^p = \ln 3 \Rightarrow p \ln 2 = \ln 3 \Rightarrow p = \frac{\ln 3}{\ln 2}$

Substituting  $2^p = 3$  into equation (1) gives,  $q = -\frac{1}{2}$

8 a  $f(x) = \frac{d}{dx} F(x)$


So where  $F(x)$  is constant,  $f(x) = 0$

$$\text{For } 1 \leq x \leq 2, f(x) = \frac{d}{dx} \frac{1}{2}(x^3 - 2x^2 + x) = \frac{3x^2}{2} - 2x + \frac{1}{2}$$

So the probability density function is:

$$f(x) = \begin{cases} \frac{3x^2}{2} - 2x + \frac{1}{2} & 1 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

b Between (1, 0) and (2, 2.5) is an arc of a positive quadratic, otherwise the function lies on the  $x$ -axis:



c  $P(X < 1.5) = F(1.5) = \frac{1}{2}(1.5^3 - 2(1.5^2) + 1.5)$

$$= \frac{1}{2} \left( \frac{27}{8} - \frac{9}{2} + \frac{3}{2} \right) = \frac{1}{2} \times \frac{3}{8} = \frac{3}{16} = 0.1875$$

9 a As area under curve must be 1,  $\int_0^2 k(4-x^2)dx = \left[ k \left( 4x - \frac{x^3}{3} \right) \right]_0^2 = 1$

$$\Rightarrow k \left( 8 - \frac{8}{3} \right) = \frac{16k}{3} = 1$$

$$\Rightarrow k = \frac{3}{16}$$

**b Method 1**

If  $x < 0$ ,  $F(x) = 0$       so  $F(0) = 0$

If  $0 \leq x \leq 2$

$$F(x) = \int_0^x \frac{3}{16}(4-t^2)dt = \left[ \frac{3}{16} \left( 4t - \frac{t^3}{3} \right) \right]_0^x = \frac{3}{16} \left( 4x - \frac{x^3}{3} \right)$$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 0 \\ \frac{3}{16} \left( 4x - \frac{x^3}{3} \right) & 0 \leq x \leq 2 \\ 1 & x > 2 \end{cases}$$

**Method 2**

If  $0 \leq x \leq 2$

$$F(x) = \int \frac{3}{16}(4-x^2) dx = \frac{3}{16} \left( 4x - \frac{x^3}{3} \right) + c$$

$$F(2) = 1 \Rightarrow \frac{3}{16} \left( 8 - \frac{8}{3} \right) + c = 1 \Rightarrow c = 0$$

This leads to the same full solution as given for Method 1.

c  $P(0.69 < X < 0.70) = F(0.70) - F(0.69) = \frac{3}{16} \left( 2.8 - \frac{0.343}{3} \right) - \frac{3}{16} \left( 2.76 - \frac{0.328509}{3} \right)$   
 $= 0.50356 - 0.49697 = 0.00659 = 0.007$  (1 s.f.)

10 a As  $F(x)$  is a cumulative distribution function,  $F(0) = 0$  and  $F(3) = 1$ . So from  $F(3) = 1$ :

$$\frac{1}{120}(3k - 27) = 1 \Rightarrow 3k = 120 + 27 = 147$$

$$\Rightarrow k = 49$$

b  $P(X > 2) = 1 - P(X \leq 2) = 1 - \frac{1}{120}(49 \times 2 - 2^3) = \frac{98 - 8}{120} = \frac{90}{120} = 0.25$

11 If  $x < 1$ ,  $F(x) = 0$  so  $F(0) = 0$

If  $1 \leq x < 7$

$$F(x) = \int \frac{1}{x \ln 7} dx = \frac{\ln x}{\ln 7} + c$$

As  $F(7) = 1$ ,  $c = 0$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 1 \\ \frac{\ln x}{\ln 7} & 1 \leq x \leq 7 \\ 1 & x > 7 \end{cases}$$

12 If  $x < 0$ ,  $F(x) = 0$  so  $F(0) = 0$

If  $0 \leq x < 0.5$

$$F(x) = \int \pi \cos(\pi x) dx = \sin(\pi x) + c$$

As  $F(0) = 0$ ,  $c = 0$

So the full solution is:

$$F(x) = \begin{cases} 0 & x < 0 \\ \sin(\pi x) & 0 \leq x \leq 0.5 \\ 1 & x > 0.5 \end{cases}$$

13 a As  $F(x)$  is a cumulative distribution function,  $F(0) = 0$  and  $F(3) = 1$ . So from  $F(3) = 1$ :

$$k(2 + \ln 3) = 1 \Rightarrow k = \frac{1}{2 + \ln 3}$$

b  $f(x) = \frac{d}{dx} F(x)$

So where  $F(x)$  is constant,  $f(x) = 0$

$$\text{For } 1 \leq x \leq 3, f(x) = \frac{d}{dx} \frac{1}{2 + \ln 3} (x - 1 + \ln x) = \frac{1}{2 + \ln 3} \left( 1 + \frac{1}{x} \right)$$

So the probability density function is:

$$f(x) = \begin{cases} \frac{1}{2 + \ln 3} \left( 1 + \frac{1}{x} \right) & 1 \leq x \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

**Challenge**

$$\mathbf{a} \quad F(t) = \int 1.25e^{-1.25t} dt = -e^{-1.25t} + c$$

$$F(0) = 0 \Rightarrow -e^0 + c = 0 \Rightarrow c = 1$$

So the full solution is:

$$F(t) = \begin{cases} 0 & t < 0 \\ 1 - e^{-1.25t} & t \geq 0 \end{cases}$$

$$\mathbf{b} \quad P(1 < T < 2) = P(T < 2) - P(T < 1) = 1 - e^{-2.5} - (1 - e^{-1.25}) = e^{-1.25} - e^{-2.5} = 0.2044 \text{ (4 d.p.)}$$

$$\mathbf{c} \quad P(T > 3) = 1 - P(T \leq 3) = 1 - (1 - e^{-3.75}) = e^{-3.75} = 0.0235 \text{ (4 d.p.)}$$