

Exercise 1D

1 a RHS = $2 \sinh A \cosh A$

$$\begin{aligned}
 &= 2 \left(\frac{e^A - e^{-A}}{2} \right) \left(\frac{e^A + e^{-A}}{2} \right) \\
 &= \frac{1}{2} (e^{2A} - 1 + 1 - e^{-2A}) \\
 &= \frac{e^{2A} - e^{-2A}}{2} \\
 &= \sinh 2A = \text{LHS}
 \end{aligned}$$

b RHS = $\cosh A \cosh B - \sinh A \sinh B$

$$\begin{aligned}
 &= \left(\frac{e^A + e^{-A}}{2} \right) \left(\frac{e^B + e^{-B}}{2} \right) - \left(\frac{e^A - e^{-A}}{2} \right) \left(\frac{e^B - e^{-B}}{2} \right) \\
 &= \frac{e^{A+B} + e^{-A+B} + e^{A-B} + e^{-A-B}}{4} \\
 &\quad - \frac{e^{A+B} - e^{-A+B} - e^{A-B} + e^{-A-B}}{4} \\
 &= \frac{2(e^{-A+B} + e^{A-B})}{4} \\
 &= \frac{e^{A-B} + e^{-(A-B)}}{2} \\
 &= \cosh(A - B) = \text{LHS}
 \end{aligned}$$

c RHS = $4 \cosh^3 A - 3 \cosh A$

$$\begin{aligned}
 &= 4 \left(\frac{e^A + e^{-A}}{2} \right)^3 - 3 \left(\frac{e^A + e^{-A}}{2} \right) \\
 (e^A + e^{-A})^3 &= e^{3A} + 3e^{2A}e^{-A} + 3e^Ae^{-2A} + e^{-3A} \\
 &= e^{3A} + 3e^A + 3e^{-A} + e^{-3A} \\
 \text{RHS} &= \frac{e^{3A} + 3e^A + 3e^{-A} + e^{-3A}}{2} - \frac{3(e^A + e^{-A})}{2} \\
 &= \frac{e^{3A} + e^{-3A}}{2} \\
 &= \cosh 3A = \text{LHS}
 \end{aligned}$$

Use the expansion
 $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$.

$$\begin{aligned}
 1 \quad d \quad \text{RHS} &= 2 \sin\left(\frac{A-B}{2}\right) \cosh\left(\frac{A+B}{2}\right) \\
 &= 2 \left(\frac{e^{\frac{A-B}{2}} - e^{-\frac{A+B}{2}}}{2} \right) \left(\frac{e^{\frac{A+B}{2}} + e^{-\frac{A-B}{2}}}{2} \right) \\
 &= \frac{1}{2} \left(e^{\frac{A-B}{2} + \frac{A+B}{2}} - e^{\frac{-A+B}{2} + \frac{A+B}{2}} + e^{\frac{A-B}{2} + \frac{A-B}{2}} - e^{\frac{-A+B}{2} + \frac{-A-B}{2}} \right) \\
 &= \frac{1}{2} (e^A - e^B + e^{-B} - e^{-A}) \\
 &= \frac{1}{2} (e^A - e^{-A}) - \frac{1}{2} (e^B - e^{-B}) \\
 &= \sinh A - \sinh B \\
 &= \text{LHS}
 \end{aligned}$$

$$\begin{aligned}
 2 \quad a \quad \sin(A-B) &= \sin A \cos B - \cos A \sin B \\
 \sinh(A-B) &= \sinh A \cosh B - \cosh A \sinh B
 \end{aligned}$$

Replace $\sin x$ by $\sinh x$ and $\cos x$ by $\cosh x$.

$$\begin{aligned}
 b \quad \sin 3A &= 3 \sin A - 4 \sin^3 A \\
 &= 3 \sin A - 4 \sin A \sin^2 A \\
 \sinh 3A &= 3 \sinh A + 4 \sinh^3 A
 \end{aligned}$$

Replace $\sin^2 A$, the product of two sine terms, by $-\sinh^2 A$.

$$\begin{aligned}
 c \quad \cos A + \cos B &= 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) \\
 \cosh A + \cosh B &= 2 \cosh\left(\frac{A+B}{2}\right) \cosh\left(\frac{A-B}{2}\right)
 \end{aligned}$$

Replace $\cos x$ by $\cosh x$.

$$\begin{aligned}
 d \quad \cos 2A &= \frac{1 - \tan^2 A}{1 + \tan^2 A} \\
 \cosh 2A &= \frac{1 + \tanh^2 A}{1 - \tanh^2 A}
 \end{aligned}$$

$\tan^2 A = \frac{\sin^2 A}{\cos^2 A}$, so there is a product of two sines.
Replace $\tan^2 A$ by $-\tanh^2 A$.

$$\begin{aligned}
 e \quad \cos 2A &= \cos^4 A - \sin^4 A \\
 &= \cos^4 A - (\sin^2 A)(\sin^2 A) \\
 \cosh 2A &= \cosh^4 A - (-\sinh^2 A)(-\sinh^2 A) \\
 &= \cosh^4 A - \sinh^4 A
 \end{aligned}$$

Replace $\sin^2 A$ by $-\sinh^2 A$.

Further Pure Maths 3

Solution Bank

3 a Using $\cosh^2 x - \sinh^2 x = 1$

$$4 - \sinh^2 x = 1$$

$$\sinh^2 x = 3$$

$$\sinh x = \pm\sqrt{3}$$

Both positive and negative values of $\sinh x$ are possible.

b Using $\tanh x = \frac{\sinh x}{\cosh x}$

$$\tanh x = \pm \frac{\sqrt{3}}{2}$$

c Using $\cosh 2x = 2\cosh^2 x - 1$

$$\cosh 2x = (2 \times 4) - 1$$

$$= 7$$

4 a Using $\cosh^2 x - \sinh^2 x = 1$

$$\cosh^2 x - (-1)^2 = 1$$

$$\cosh^2 x = 2$$

$$\cosh x = \sqrt{2}$$

$\cosh x$ cannot be negative.

b Using $\sinh 2x = 2\sinh x \cosh x$

$$\sinh 2x = 2 \times (-1) \times \sqrt{2}$$

$$= -2\sqrt{2}$$

c Using $\tanh 2x = \frac{2 \tanh x}{1 + \tanh^2 x}$

$$\tanh x = \frac{\sinh x}{\cosh x} = \frac{-1}{\sqrt{2}}$$

$$\tanh 2x = \frac{\left(-\frac{2}{\sqrt{2}}\right)}{1 + \left(\frac{1}{2}\right)}$$

$$= \frac{-2}{\sqrt{2}} \times \frac{2}{3}$$

$$= \frac{-4}{3\sqrt{2}} = -\frac{2\sqrt{2}}{3}$$

Alternatively use
 $\frac{\sinh 2x}{\cosh 2x} = \frac{\sinh 2x}{2\cosh^2 x - 1}$

5 a

$$3\sinh x + 4\cosh x = 4$$

$$\frac{3(e^x - e^{-x})}{2} + \frac{4(e^x + e^{-x})}{2} = 4$$

$$3e^x - 3e^{-x} + 4e^x + 4e^{-x} = 8$$

$$7e^x - 8 + e^{-x} = 0$$

$$7e^{2x} - 8e^x + 1 = 0$$

$$(7e^x - 1)(e^x - 1) = 0$$

$$e^x = \frac{1}{7} \text{ or } e^x = 1$$

$$x = \ln\left(\frac{1}{7}\right), x = 0$$

Multiply throughout by e^x .

Solve as a quadratic in e^x .

Note that
 $\ln\left(\frac{1}{7}\right) = \ln(7^{-1})$
 $= -\ln 7$

b

$$7\sinh x - 5\cosh x = 1$$

$$\frac{7(e^x - e^{-x})}{2} - \frac{5(e^x + e^{-x})}{2} = 1$$

$$7e^x - 7e^{-x} - 5e^x - 5e^{-x} = 2$$

$$2e^x - 2 - 12e^{-x} = 0$$

$$e^x - 1 - 6e^{-x} = 0$$

$$e^{2x} - e^x - 6 = 0$$

$$(e^x - 3)(e^x + 2) = 0$$

$$e^x = 3$$

$$x = \ln 3$$

Multiply throughout by e^x .

$e^x = -2$ is not possible for real x .

c

$$30\cosh x = 15 + 26\sinh x$$

$$30\frac{(e^x + e^{-x})}{2} = 15 + 26\frac{(e^x - e^{-x})}{2}$$

$$15e^x + 15e^{-x} = 15 + 13e^x - 13e^{-x}$$

$$2e^x - 15 + 28e^{-x} = 0$$

$$2e^{2x} - 15e^x + 28 = 0$$

$$(2e^x - 7)(e^x - 4) = 0$$

$$e^x = \frac{7}{2}, e^x = 4$$

$$x = \ln\left(\frac{7}{2}\right), x = \ln 4$$

Multiply throughout by e^x .

Solve as a quadratic in e^x .

5 d $13 \sinh x - 7 \cosh x + 1 = 0$

$$13 \frac{(e^x - e^{-x})}{2} - 7 \frac{(e^x + e^{-x})}{2} + 1 = 0$$

$$13e^x - 13e^{-x} - 7e^x - 7e^{-x} + 2 = 0$$

$$6e^x + 2 - 20e^{-x} = 0$$

$$3e^x + 1 - 10e^{-x} = 0$$

$$3e^{2x} + e^x - 10 = 0$$

$$(3e^x - 5)(e^x + 2) = 0$$

$$e^x = \frac{5}{3}$$

$$x = \ln\left(\frac{5}{3}\right)$$

Multiply throughout by e^x .

Solve as a quadratic in e^x .

$e^x = -2$ is not possible for real x .

e

Use $\operatorname{arsinh} x = \ln(x + \sqrt{x^2 + 1})$.

$$\cosh 2x - 5 \sinh x = 13$$

Using $\cosh 2x = 1 + 2 \sinh^2 x$,

$$1 + 2 \sinh^2 x - 5 \sinh x = 13$$

$$2 \sinh^2 x - 5 \sinh x - 12 = 0$$

$$(2 \sinh x + 3)(\sinh x - 4) = 0$$

$$\sinh x = -\frac{3}{2}, \sinh x = 4$$

$$x = \operatorname{arsinh}\left(-\frac{3}{2}\right), x = \operatorname{arsinh} 4$$

$$x = \ln\left(-\frac{3}{2} + \sqrt{\frac{9}{4} + 1}\right)$$

$$= \ln\left(\frac{-3 + \sqrt{13}}{2}\right)$$

$$x = \ln(4 + \sqrt{16 + 1})$$

$$= \ln(4 + \sqrt{17})$$

5 f $3\sinh^2 x - 13\cosh x + 7 = 0$

Using $\cosh^2 x - \sinh^2 x = 1$,

$$3(\cosh^2 x - 1) - 13\cosh x + 7 = 0$$

$$3\cosh^2 x - 13\cosh x + 4 = 0$$

$$(3\cosh - 1)(\cosh x - 4) = 0$$

$$\cosh x = \frac{1}{3}, \cosh x = 4$$

$$\cosh x = 4$$

$$x = \operatorname{arcosh} 4, -\operatorname{arcosh} 4$$

$$x = \ln(4 \pm \sqrt{4^2 - 1})$$

$$= \ln(4 \pm \sqrt{15})$$

$\cosh x \geq 1$, so $\cosh x = \frac{1}{3}$ is not possible.

Use $\operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$, but remember that $\ln(x - \sqrt{x^2 - 1})$ is also a solution.

g $\sinh 2x - 7\sinh x = 0$

$$2\sinh x \cosh x - 7\sinh x = 0$$

$$\sinh x(2\cosh x - 7) = 0$$

$$\sinh x = 0, \cosh x = \frac{7}{2}$$

$$x = 0, x = \pm \operatorname{arcosh}\left(\frac{7}{2}\right)$$

$$\operatorname{arcosh}\left(\frac{7}{2}\right) = \ln\left(\frac{7}{2} + \sqrt{\frac{49}{4} - 1}\right)$$

$$= \ln\left(\frac{7 + \sqrt{45}}{2}\right)$$

$$= \ln\left(\frac{7 + 3\sqrt{5}}{2}\right)$$

$$x = 0, x = \ln\left(\frac{7 \pm 3\sqrt{5}}{2}\right)$$

Use $\operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$, but remember that $\ln(x - \sqrt{x^2 - 1})$ is also a solution.

h $4\cosh x + 13e^{-x} = 11$

$$4\frac{(e^x + e^{-x})}{2} + 13e^{-x} = 11$$

$$2e^x + 2e^{-x} + 13e^{-x} = 11$$

$$2e^x + 15e^{-x} - 11 = 0$$

$$2e^{2x} - 11e^x + 15 = 0$$

$$(2e^x - 5)(e^x - 3) = 0$$

$$e^x = \frac{5}{2}, e^x = 3$$

$$x = \ln\left(\frac{5}{2}\right), x = \ln 3$$

Multiply throughout by e^x .

Solve as a quadratic in e^x .

5 i

$$2 \tanh x = \cosh x$$

$$\frac{2 \sinh x}{\cosh x} = \cosh x$$

$$2 \sinh x = \cosh^2 x$$

Using $\cosh^2 x - \sinh^2 x = 1$

$$2 \sinh x = 1 + \sinh^2 x$$

$$\sinh^2 x - 2 \sinh x + 1 = 0$$

$$(\sinh x - 1)^2 = 0$$

$$\sinh x = 1$$

$$x = \operatorname{arsinh} 1$$

$$x = \ln(1 + \sqrt{1^2 + 1})$$

$$= \ln(1 + \sqrt{2})$$

Use $\operatorname{arsinh} x = \ln(x + \sqrt{x^2 + 1})$.

6 a

$$\begin{aligned} 2 \cosh^2 x - 1 &= 2 \left(\frac{e^x + e^{-x}}{2} \right)^2 - 1 \\ &= 2 \left(\frac{e^{2x} + 2 + e^{-2x}}{4} \right) - 1 \\ &= \frac{e^{2x} + e^{-2x}}{2} \\ &= \cosh 2x. \end{aligned}$$

b Using the relation $\cosh 2x = 2 \cosh^2 x - 1$, the equation becomes $2 \cosh^2 x - 3 \cosh x - 9 = 0$. Substituting $u = \cosh x$ into this equation allows us to clearly see a quadratic equation

$2u^2 - 3u - 9 = 0$ which gives solutions $u = \frac{3 \pm \sqrt{9 + 72}}{4}$, we neglect the negative solution to avoid

complex numbers.

i.e.

$$u = 3.$$

This means that our solutions for x are the values for

$$x_1 = \operatorname{ar} \cosh(3).$$

Now recalling the expression $\operatorname{ar} \cosh u = \ln(u \pm \sqrt{u^2 - 1})$, we obtain the exact logarithm solutions

$$x = \ln(3 \pm 2\sqrt{2}).$$

7 Using the identity $\cosh^2 x - \sinh^2 x \equiv 1$, the equation in question becomes

$$2(\cosh^2 x - 1) - 5 \cosh x = 5.$$

We simplify and use the substitution $u = \cosh x$ in order to obtain $2u^2 - 5u - 7 = 0$. This equation has

solutions $u = \frac{5 \pm \sqrt{25 + 56}}{4}$, we neglect the negative solution to avoid complex numbers, thus $u = \frac{7}{2}$.

This means that our solutions for x are the values for

$$x = \operatorname{arccosh}\left(\frac{7}{2}\right).$$

Now recalling the expression $\operatorname{arccosh} u = \ln\left(u \pm \sqrt{u^2 - 1}\right)$, we obtain the exact logarithm solutions

$$x = \ln\left(\frac{1}{2}(7 \pm 3\sqrt{5})\right).$$

8 $\operatorname{sech}^2 x \equiv 1 + \tanh^2 x$ is not true, the correct identity is $\operatorname{sech}^2 x \equiv 1 - \tanh^2 x$ by Osborn's rule. He has split the fraction via denominator on the second line. This is not valid.

In mathematics we write this as

$$\frac{a}{b+c} \neq \frac{a}{b} + \frac{a}{c}.$$

The final mistake he made was assuming that taking the reciprocal of all terms preserves the value. This is incorrect. For example $3+1 \neq \frac{1}{3}+1$.

The correct proof is

$$\begin{aligned} \frac{1 + \tanh^2 x}{1 - \tanh^2 x} &\equiv \frac{2 - \operatorname{sech}^2 x}{\operatorname{sech}^2 x} \\ &\equiv \frac{2}{\operatorname{sech}^2 x} - 1 \\ &\equiv 2 \cosh^2 x - 1. \end{aligned}$$

9 a Recall the identity $R \cosh(x+a) \equiv R \cosh x \cosh a + R \sinh x \sinh a$.

From this we can set $R \cosh a = 10$ and $R \sinh a = 6$.

In order to find values for R and a :

$$\frac{R \sinh a}{R \cosh a} = \tanh a = \frac{6}{10} = \frac{3}{5}$$

giving

$$\begin{aligned} a &= \operatorname{artanh}(0.6) \\ &\approx 0.693(3 \text{ d.p.}) \end{aligned}$$

We can now use the expression $R \cosh a = 10$.

$$\text{To find } \cosh a : \operatorname{sech}^2 a = 1 - \tanh^2 a = 1 - \left(\frac{3}{5}\right)^2 = \frac{16}{25}$$

$$\therefore \cosh^2 a = \frac{1}{\operatorname{sech}^2 a} = \frac{25}{16} \text{ and so } \cosh a = \frac{5}{4}$$

$$\text{So } R = \frac{10}{\cosh a} = \frac{10}{\left(\frac{5}{4}\right)} = 8$$

Thus giving us $10 \cosh x + 6 \sinh x = 8 \cosh(x + 0.693)$.

- 9 b** The minimum value occurs when $\cosh(x+0.693)$ is minimal. We know this occurs when $x+0.693=0$, by the graph of $\cosh x$. So the minimal value is $8\cosh 0=8$.

We could also find this answer by setting $y=8\cosh(x+0.693)$, finding the solution to

$$\frac{dy}{dx}=8\sinh(x+0.693)=0 \text{ as } x=-0.693 \text{ and substituting in, to get } y=8\cosh 0=8.$$

The second derivative should be checked to be positive at this point in order to conclude that the stationary point is minimal, not maximal.

- c** $10\cosh x+6\sinh x=8\cosh(x+0.693)=11$.

We rearrange the equation to be $\cosh(x+0.693)=\frac{11}{8}$.

Set $u=x+0.693$ and now noting the symmetry about $u=0$ we obtain two solutions,

$$u=\pm\operatorname{arcosh}\left(\frac{11}{8}\right)\approx\pm 0.841(3 \text{ d.p.})$$

This means our solutions for x are

$$x_1=-1.534,$$

$$x_2=0.148.$$